Suczt Dro[®] (114 kg.) Net Wt. 250 lbs.

Minimum 18.00%

Guaranteed Analysis

Crude Fat	00%
Crude Fiber	50%
Calcium (Ca) Minimum 1.75% - Maximum 2.2	25%
Phosphorus (P)	
Salt (NaCl)	
Magnesium (Mg)	25%
Potassium (K)	
Sulfur (S) Minimum 1.0	00%
Copper (Cu) Minimum 600 p	ppm
Selenium (Se)	ppm
Zinc (Zn) Minimum 1,200 p	ppm
Manganese (Mn)	ppm
Cobalt (Co) Minimum 4.0 p	ppm
Iodine (I) Minimum 8.0 p	
Vitamin A	U/lb.
Vitamin D	U/lb.
Vitamin E	U/lb.
Alpha-Amylase (from Aspergillus oryzae)	

Ingredients: Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Salt, Magnesium Oxide, Mono-Calcium Phosphate, Corn Gluten Meal, Calcium Carbonate, Calcium Sulfate, Canola Oil, Biuret, Ammonium Sulfate, Potassium Chloride, Yeast Culture, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Dried Aspergillus Oryzae Fermentation Extract, Liquid Aspergillus Niger Fermentation Product, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

FiberMate™ 18

SweetPro® FiberMate 18 lick blocks are designed for essential fortification of Yearling Cattle and Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. FiberMate 18 should be used for cattle on average to below average forages. Daily intake will vary depending on forage quality and nutritional needs.

Feeding Instructions:

- Before feeding SweetPro® FiberMate 18 lick blocks for the first time, animals should be well fed.
- 2. Provide access for animals to have plenty of clean, fresh water at all times.
- 3. Provide an adequate source of good quality roughage and forage at all times (all they can consume).
- 4. Have one lick block available per 25 head of beef or dairy cattle.
- 5. Place lick blocks near water sources or areas cattle frequent regularly.
- Make lick blocks available in more than one location in large pastures or where more than one water source is available.
- 7. Other salt and minerals may be provided free choice to balance specific needs.
- 8. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .75 to 1.25 lbs./hd/day.
- Once animals consistently consume more than 1.25 lbs./hd/day, transition to FiberMate 20. If daily consumption is consistently below .75 lbs./hd/day, transition to SweetPro® 16.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

Caution!

Contains 600 ppm copper. **DO NOT FEED TO SHEEP!**Do not use in association with another feed containing supplemental selenium.

Control #:_____



Crude Protein