SWZZLDCO® (114 kg.) Net Wt. 250 lbs.

Guaranteed Analysis

	Minimum 30.00%
(includes not more than 15% equi-	valent protein from non-protein-nitrogen)
Crude Fat	
Calcium (Ca)	
Phosphorus (P)	Minimum 1.40%
Salt (NaCl)	
Magnesium (Mg)	
Potassium (K)	Minimum 1.00%
Copper (Cu)	Minimum 600 ppm
Selenium (Se)	
Zinc (Zn)	Minimum 1,200 ppm
Manganese (Mn)	
	Minimum 4.0 ppm
Iodine (I)	
Vitamin A	Minimum 100,000 IU/lb.
Vitamin D	
Amyloglucosidase (from Aspergillus niger, Aspergill	us oryzae)Minimum 513 ug maltose hydrolyzed/min/oz.

Ingredients: Corn Distillers Dried Grains with Solubles, Corn Condensed Distillers Solubles, Salt, Corn Gluten Meal, Ammonium Sulfate, Magnesium Oxide, Mono-Calcium Phosphate, Biuret, Calcium Carbonate, Calcium Sulfate, Potassium Chloride, Yeast Culture, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Aspergillus Oryzae Extract, Aspergillus Niger Fermentation Liquid, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

Magnum™

SweetPro* MagnumTM tubs are designed for essential fortification of Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. Magnum should be used for cattle on below average to poor forages. Daily intake will vary depending on forage quality and nutritional needs.

Feeding Instructions:

- Before feeding SweetPro® Magnum lick tubs for the first time, animals should be well fed.
- 2. Provide access for animals to have plenty of clean, fresh water at all times.
- Provide an adequate source of good quality roughage and forage at all times (all they can consume).
- 4. Have one lick tub available per 25 head of beef cattle.
- 5. Place tubs near water sources or areas cattle frequent regularly.
- 6. Make tubs available in more than one location in large pastures or where more than one water source is available.
- 7. Other salt and minerals may be provided free choice to balance specific needs.
- A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .75 to 1.25 lbs. per day.
- Once animals consistently consume less than .75 lbs./hd/day, transition to SweetPro® Super 25.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

Caution!

Contains 600 ppm copper. DO NOT FEED TO SHEEP!

Do not use in association with another feed containing supplemental selenium. This feed shall be used only in accordance with directions furnished on the label.

Control #:

