SUCZCLDICO® (102 kg.) Net Wt. 225 lbs.

Minimum 16 000/

Guaranteed Analysis

Crude Protein	Minimum 16.00%
Crude Fat	Minimum 5.00%
Crude Fiber	
Calcium (Ca)	
Phosphorus (P)	
Salt (NaCl)	
	Minimum 1.00%
Potassium (K)	
Copper (Cu)	Minimum 600 ppm
Selenium (Se)	
Zinc (Zn)	Minimum 1,200 ppm
Manganese (Mn)	
Cobalt (Co)	
Iodine (I)	Minimum 8.0 ppm
Vitamin A	Minimum 100,000 IU/lb.
	Minimum 3,393 ug starch hydrolyzed/min/oz.
Amyloglucosidase (from Aspergillus niger, Aspergillus	s oryzae)Minimum 513 ug maltose hydrolyzed/min/oz.

Ingredients: Com Distillers Dried Grains with Solubles, Com Condensed Distillers Solubles, Salt, Sodium Bicarbonate, Mono-Calcium Phosphate, Calcium Carbonate, Magnesium Oxide, Potassium Chloride, Yeast Culture, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, DriedAspergillus Oryzae Fermentation Extract, Liquid Aspergillus Niger Fermentation Product, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

Starter

SweetPro® Starter lick blocks are designed for essential fortification of Receiving Calves and Stocker Cattle requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. Daily intake will vary depending on forage quality and nutritional needs.

Feeding Instructions:

- 1. Have one lick block available per 15-25 head of beef or dairy cattle.
- 2. Provide an adequate source of good quality roughage or forage at all times (all they can consume).
- 3. Provide access to plenty of clean fresh water at all times.
- 4. When starting Calves, place SweetPro® Starter lick blocks close to water sources or in areas where animals will have frequent access to the product.
- For Stocker Cattle on good to excellent pasture, make lick blocks available in more than one location in large pastures or where more than one water source is available.
- 6. Other salt and minerals may be provided free choice to balance specific needs.
- 7. Recommended daily intake of product is 0.75 to 1.25 lbs./hd/day.
- Once cattle consistently consume SweetPro® Starter above 1.25 lbs./hd/day, transition animals to SweetPro® Cattle Kandi.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

Caution!

Contains 600 ppm copper. **DO NOT FEED TO SHEEP!**

Do not use in association with another feed containing supplemental selenium.

Control #:_____



C------