# Sucation Net Wt. 250 lbs.

## **Guaranteed Analysis**

(includes not more than 8% equivalent protein from non-protein-nitrogen)	
Crude Fiber	
Calcium (Ca)	
Salt (NaCl)	Minimum 12.00% - Maximum 13.50%
Potassium (K)	
Sulfur (S)	Minimum 1.50%
Copper (Cu)	Minimum 600 ppm
	Minimum 5.0 ppm - Maximum 5.25 ppm
Zinc (Zn)	Minimum 1,200 ppm
Manganese (Mn)	Minimum 750 ppm
Cobalt (Co)	Minimum 4.0 ppm
Vitamin A	
Vitamin E	
Alpha-Amylase (from Aspergillus oryzae)	Minimum 3,393 ug starch hydrolyzed/min/oz.
Amyloglucosidase (from Aspergillus niger, Aspergillus	oryzae)Minimum 513 ug maltose hydrolyzed/min/oz.

Ingredients: Com Distillers Dried Grains with Solubles, Com Condensed Distillers Solubles, Salt, Magnesium Oxide, Com Gluten Meal, Mono-Calcium Phosphate, Ammonium Sulfate, Calcium Carbonate, Vegetable Oil, Biuret, Calcium Sulfate, Potassium Chloride, Yeast Culture, Zinc Sulfate, Manganese Sulfate, Copper Sulfate, Zinc Polysaccharide Complex, Zinc Proteinate, Manganese Polysaccharide Complex, Manganese Proteinate, Copper Polysaccharide Complex, Copper Proteinate, Dried Aspergillus Oryzae Fermentation Extract, Liquid Aspergillus Niger Fermentation Product, Cobalt Carbonate, Sodium Selenite, Ethylenediamine Dihydroiodide, Vitamin A Acetate (Stability Improved), Vitamin D3 Supplement, Vitamin E Supplement.

Zinc, Manganese and Copper are 50% organically complexed.

Does Not Contain Any Animal Protein

# Super 25

SweetPro® Super 25 lick blocks are designed for the essential fortification of Cow requirements for vitamins A, D & E, plus protein, calcium, phosphorus and trace elements including selenium. Super 25 should be used for cattle on below average to poor forages. Daily intake will vary depending on forage quality and nutritional needs.

# Feeding Instructions:

- Before feeding SweetPro® Super 25 lick blocks for the first time, animals should be well fed.
- 2. Provide access for animals to have plenty of clean, fresh water at all times.
- 3. Have one lick block available per 25 head of beef cows.
- Provide an adequate source of good quality roughage and forage at all times (all they can consume).
- 5. Place lick blocks near water sources or areas cattle frequent regularly.
- Make lick blocks available in more than one location in large pastures or where more than one water source is available.
- 7. Other salt and minerals may be provided free choice to balance specific needs.
- 8. A two week adaptation period will be required to establish a base consumption rate. After adaptation period, recommended range of intake is .75 to 1.25 lbs. per day.
- Once animals consistently consume more than 1.25 lbs/hd/day, transition to SweetPro Magnum. If daily consumption is consistently below .75 lbs/hd/day, transition to SweetPro FiberMate 20.

The above feeding directions are general guidelines and will vary with the overall feeding conditions. Always use good management practices.

# Caution!

Contains 600 ppm copper. DO NOT FEED TO SHEEP!

Do not use in association with another feed containing supplemental selenium. This feed shall be used only in accordance with directions furnished on the label.

