



Figure 1

Example of a current feed grade antibiotic label

Generic CTC (Chlortetracycline) 100 NADA 48-480

Generic Name: chlortetracycline hydrochloride granule Dosage Form: FOR ANIMAL USE ONLY

CALVES

DOSAGE LEVEL

0.1 mg/lb. body wt daily

INDICATIONS FOR USE

Calves (up to 250 lbs.): For an increased rate of body weight gain and improved feed efficiency.

25-70 mg/head/day

Calves (250-400 lbs.) For an increased rate of weight gain and improved feed efficiency

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

GROWING CATTLE

DOSAGE LEVEL

70 mg/head/day

INDICATION FOR USE

Growing cattle (over 400 lbs.): For an increased rate of weight gain, improved feed efficiency and for reduction of incidence of liver abscesses in beef cattle associated with Fusobacterium necrophorum and Arcanobacterium pyogenes

WARNING: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

BEEF CATTLE

DOSAGE LEVEL

350 mg/head/day

INDICATION FOR USE

For the control of bacterial pneumonia associated with shipping fever complex caused by Pasteurella spp. susceptible to chlortetracycline.

350 mg/head/day

Beef cattle (under 700 lbs.): Control of active infection of anaplasmosis caused by Anaplasma marginale susceptible to chlortetracycline.

0.5 mg/lb. daily

Beef cattle (Over 700 lbs.): Control of active infection of anaplasmosis caused by Anaplasma marginale susceptible to chlortetracycline.

WARNING: Withdraw 48 hours prior to slaughter.

FOR CALVES, BEEF AND NON-LACTATING DAIRY CATTLE

DOSAGE LEVEL

10 mg/lb. body wt daily

INDICATION FOR USE

Treatment of bacterial enteritis caused by Escherichia coli and bacterial pneumonia caused by Pasteurella multocida organisms susceptible to chlortetracycline. (Treat for not more than 5 days. Feed approximately 400 g/ton, varying with body weight and feed consumption to provide 10 mg/lb. body weight per day.)

WARNING: Withdraw 10 days prior to slaughter. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.